

Question

Kota Saito

January 19, 2022

Let X be the set of alternatives. Let $\mathcal{D} \subset 2^X \setminus \emptyset$ be the set of choice sets.

Stochastic choice function: A function $\rho : \mathcal{D} \times X \rightarrow [0, 1]$ is called a *stochastic choice function* if $\sum_{x \in D} \rho(D, x) = 1$ and $\rho(D, x) = 0$ for any $x \notin D$.

For each $(D, x) \in \mathcal{D} \times X$, the number $\rho(D, x)$ is the probability that an alternative x is chosen from a choice set D .

Rankings: Let Π be the set of bijections between X and $\{1, \dots, |X|\}$, where $|X|$ is the number of elements of X . If $\pi(x) = i$, then we interpret x to be the $|X| + 1 - i$ -th best element of X with respect to π . If $\pi(x) > \pi(y)$, then x is better than y with respect to π . An element π of Π is called a *strict preference ranking* (or simply, a *ranking*) over X . For all $(D, x) \in \mathcal{D} \times X$ such that $x \in D$, if $\pi(x) > \pi(y)$ for all $y \in D \setminus \{x\}$, then we often write $\pi(x) \geq \pi(D)$. There are $|X|!$ elements in Π .

$$\rho^\pi(D, x) = \begin{cases} 1 & \text{if } \pi(x) \geq \pi(D); \\ 0 & \text{otherwise.} \end{cases} \quad (1)$$

The function ρ^π gives probability one to the best alternative x in a choice set D according to the strict preference ranking π . The following fact is elementary but fundamental:

We denote the set of probability measures over Π by $\Delta(\Pi)$. Since Π is finite, $\Delta(\Pi) = \{(\nu_1, \dots, \nu_{|\Pi|}) \in \mathbf{R}_+^{|\Pi|} \mid \sum_{i=1}^{|\Pi|} \nu_i = 1\}$, where \mathbf{R}_+ is the set of nonnegative real numbers.

We now introduce the definition of random utility model:

Definition 1. A stochastic choice function ρ is called a *random utility function* if there

22 exists a probability measure $\nu \in \Delta(\Pi)$ such that for all $(D, x) \in \mathcal{D} \times X$, if $x \in D$, then

$$\rho(D, x) = \nu(\pi \in \Pi | \pi(x) \geq \pi(D)).$$

23 The probability measure ν is said to *rationalize* ρ . The set of random utility functions is
 24 denoted by \mathcal{P}_r .¹

25 **Observation:** *The set \mathcal{P}_r of random utility functions is a polytope, that is $\mathcal{P}_r = \text{co.}\{\rho^\pi | \pi \in$
 26 $\Pi\}$.*

27 The observation holds because any random utility function $\rho \in \mathcal{P}_r$ can be written as
 28 $\rho = \sum_{\pi \in \Pi} \nu(\pi) \rho^\pi$ for some probability measure ν , which is guaranteed to exist by the
 29 definition of a random utility function.




Figure 1: Random utility polytope

30 The hexagons in Figure 1 illustrates the polytope. Although the geometric intuition is
 31 useful, it is important to notice that the figure oversimplifies the reality since the number
 32 (i.e., $|X|!$) of vertices and the dimension of a random utility function can be very large.²

33 One concept that is useful for analyzing the polytope is *adjacency*. As you can see, in
 34 Figure 1, ρ^{π_i} and $\rho^{\pi_{i+1}}$ are *adjacent* for each $i \leq 6$.³

35 This can be formalized as follows:

36 **Definition 2.** *The two rankings π and π' are adjacent if there exists $t \in \mathbf{R}^{\mathcal{D} \times X}$ and $a \in \mathbf{R}$
 37 such that (i) $\rho^\pi \cdot t = a = \rho^{\pi'} \cdot t$ and (ii) for any $\hat{\pi}$, if $\pi \neq \hat{\pi} \neq \pi'$, then $\rho^{\hat{\pi}} \cdot t > a$.*

¹While the function above is often called a random ranking function, a random utility function is often defined differently—by using the existence of a probability measure μ over utilities such that for all $(D, x) \in \mathcal{D} \times X$, if $x \in D$, then $\rho(D, x) = \mu(u \in \mathbf{R}^X | u(x) \geq u(D))$.

²To see why the dimension of a random utility function can be very large, notice that it assigns a number for each pair of $(D, x) \in \mathcal{D} \times X$.

³Assuming $\rho^{\pi_7} = \rho^{\pi_1}$.

38 For example, take ρ^{π_1} and ρ^{π_2} . Consider a line that passes through ρ^{π_1} and ρ^{π_2} . Notice
39 that only ρ^{π_1} and ρ^{π_2} are on the line and all of the other $\{\rho^{\pi_i}\}_{i \neq 1,2}$ are below the line.
40 Thus, conditions (i) and (ii) are indeed satisfied.

41

42 **Question:** Please answer the following two questions assuming $X = \{1, 2, 3\}$:

43

44 **Question 1:** Assume $\mathcal{D} = \{\{1, 2\}, \{1, 3\}, \{2, 3\}\}$. Find out all adjacency pairs. Do your
45 best to try to find all of them.

46

47 **Question 2:** Assume $\mathcal{D} = \{\{1, 2\}, \{1, 3\}, \{2, 3\}, \{1, 2, 3\}\}$. Find out all adjacency pairs.
48 Do your best to try to find all of them.